Mike Milburn Chair Montana

Doug Grob Montana

Jeffery C. Allen Idaho

Ed Schriever Idaho



Thomas L (Les) Purce Vice Chair Washington

> KC Golden Washington

Margaret Hoffmann Oregon

Charles F. Sams III
Oregon

October 7, 2025

MEMORANDUM

TO: Council Members

FROM: Windy Schoby

SUBJECT: White sturgeon status report for the Snake River between Hells Canyon Dam

and the Lower Granite pool

BACKGROUND:

Presenter: Joe DuPont, Clearwater Regional Fisheries Manager (Idaho Department of Fish

and Game)

Summary:

The Idaho Department of Fish and Game, in collaboration with Idaho Power, is studying White Sturgeon populations in the Snake River between Hells Canyon Dam and Lower Granite Dam. The presentation will review population status, factors influencing abundance, ongoing efforts, and potential future actions to support long-term persistence.

Research indicates that juvenile sturgeon abundance in Hells Canyon has declined since the early 1970s. Juvenile growth rates are among the slowest in the Columbia Basin, reducing the number reaching adulthood. Over the past decade, meaningful recruitment has occurred only once, and the total population has declined by approximately 33%. It is estimated that more than 50% of each year-class is lost downstream of Lower Granite Dam, likely contributing to the long-term decline in juvenile abundance.

Recent changes in the biological community within the Lower Granite Reservoir appear to be affecting recruitment and survival. Trap-and-haul pilot studies, both

upstream and downstream, are underway to evaluate whether this approach can improve growth and increase spawning abundance. Additionally, the University of Idaho is developing a population model to assess long-term trends and evaluate the potential benefits of different management actions.

Relevance:

White Sturgeon are listed as an <u>emerging priority</u> in the Council's 2014 Fish and Wildlife Program and 2020 addendum. This report provides information to the region on the status and management of a portion of the Snake River.

Background: Snake River White Sturgeon Management- (IDFG 2024)

White Sturgeon have declined from historical abundances across their range, including Idaho. Historical declines are a combination of factors including overfishing, habitat manipulation, poor water quality, bioaccumulation of contaminants, and altered fish assemblages in larger river systems. However, habitat alteration and fragmentation from dam construction during 1900 – 1980 are considered the primary factors in declines in wild populations within Idaho. Dam construction and operations blocked migrations and altered flows, water temperatures, and nutrient regimes.

Two genetically distinct White Sturgeon populations exist in the Snake River:

- Lower Snake River Population (below Hells Canyon Dam) genetically similar to lower Columbia River sturgeon.
- Middle Snake River Population (upstream of Brownlee Dam) genetically distinct and fragmented by six dams.

Below Hells Canyon Dam, there is one continuous free-flowing reach along the Idaho border, but outside of Idaho, the lower Snake River genetic population is fragmented by the four lower Snake River dams in the state of Washington. Above Brownlee Dam there are six management reaches within the Middle Snake River genetic population. Because of very low abundance and limited genetic information available for Oxbow or Hells Canyon reservoirs, it is unknown at this time whether these reaches are included in the Lower Snake or Middle Snake genetic populations.

Reaches from Lower Granite Dam to Hells Canyon Dam and C.J. Strike Dam to Bliss Dam support natural spawning and recruitment of White Sturgeon previously resulting in stable or increasing abundance. Remaining reaches in the middle Snake River require hatchery stocking or downstream movement of sturgeon to support reach abundance as limited or no natural recruitment occurs in these reaches.

Some reaches, such as from Lower Granite Dam to Hells Canyon Dam and C.J. Strike to Bliss Dam, still support natural spawning and recruitment. Other reaches rely on hatchery stocking or downstream movement of fish to maintain abundance.

The White Sturgeon hatchery program, jointly operated by IDFG and Idaho Power, collects fertilized eggs from the wild (C.J. Strike–Bliss reach), hatches and rears them in a hatchery, and releases juveniles into reaches with limited recruitment—a process known as **repatriation**. The program also supports recreational angling opportunities upstream of Shoshone Falls.


Management reaches are categorized as follows:

- **Core Wild:** Self-sustaining populations with adequate natural recruitment. Management focuses on maintaining environmental conditions that support spawning and survival.
- **Stocked:** Populations dependent on hatchery supplementation due to habitat limitations such as flow or fragmentation.
- **Non-native Range:** Reaches upstream of Shoshone Falls, supported solely through hatchery stocking to provide angling opportunities.

More info:


- Idaho Department of Fish and Game Snake River White Sturgeon Management Plan 2024-2033
- 2025 Joint Staff Report Stock status and fisheries for sturgeon and smelt
- The Council's White Sturgeon web page
- Columbia Basin White Sturgeon Planning <u>Framework</u>
- White Sturgeon Story Map





Reservoir 36 miles

Free flowing River 101 miles



1972 - 1975 (Coon et al. 1977)

- 876 (204 recaps) WS captures using set lines and rod-and-reel
- Work only occurred in pools in free-flowing river

1982 - 1984 (Lukens 1985)

- 531 (49 recaps) WS captures using set lines and rod-and-reel
- Work occurred in river from Captain John to Johnson Bar

1990 - 1991 (Lepla 1994)

- 946 (147 recaps) WS captures using mostly gill nets
- •Work occurred in Lower Granite pool

1997 - 2002 (Lepla et al. 2001; Everett et al. 2002)

- 1,857 WS captures (521 recaps) using set lines
- •Work occurred from Lower Granite Dam to Granite Rapids


2012 - 2014 (Bentz 2015)

- 1,483 (281 recaps) WS captures using set lines
- 37% of catch < 3ft
- Work occurred from Lower Granite Dam to Hells Canyon Dam

2024 - 2025 (Bentz in progress)

Work occurred from Lower Granite Dam to Hells Canyon Dam





From 1960-1969 3-6 ft fish could be harvested

Starting 1970 no harvest was allowed















White Sturgeon Recruitment in Lower Granite Pool


















White Sturgeon Population Estimate in the Hells Canyon Reach



What is going on?

Why has a long-term declining trend in juvenile abundance occurred?

Why has the more recent decline in recruitment occurred?















Why has a long-term declining trend in juvenile abundance occurred?



Sturgeon Growth Rates in Hells Canyon





Mean Fork Length (10 cm increments)



Entrainment










White Sturgeon Counted in Juvenile Fish Bypass Systems (Fish Separator)







My Thoughts on Entrainment



Why the recent decline in recruitment?

Decline in recruitment (to age-1)








First showed up in Lower Granite in 1994


Now they represent over 90% of the epibenthic biomass in the reservoir (15-200 m²)


Live throughout the reservoir although prefer slower velocities

Average length: 12 mm



Siberian Praws Counted at Lower Granite Dam









Sand Rollers



Native to Columbia basin


First collected in Snake River at Lower Granite in 2003


Prefer smaller substrates

Occupy shallow and deep habitats


Main food item is opossum shrimp although larval fish were found in their diet

Sand Rollers Counted at Lower Granite Dam






Sand Roller and Sturgeon Counts and Lower Granite Dam



Sand Roller and Sturgeon Counts and Lower Granite Dam













211 fish translocated
Between 2021 to 2023
42 were acoustic
tagged







Moved 80 sturgeon


Nisqually John

- 13 acoustic tagged adults
- 8 acoustic tagged sub-adults
- 36 with only PIT tag


Heller Bar

- 14 acoustic tagged adults
- 9 acoustic tagged sub-adults










Average flow in Snake River at Anatone (May-July)







Summary

- Recruitment likely has not been sufficient to sustain the Hells Canyon White Sturgeon population since the Lower Snake River Dam Complex was completed (Lower Granite was impounded on Feb 14, 1975).
- Over 50% of each year class is believed to be entrained over Lower Granite Dam.
- Slow growth rates of juvenile sturgeon limit the number of fish that reach maturity and likely limit recruitment in the free-flowing river.
- A 33% decline is sturgeon abundance has occurred over the last 10 years in the Hells Canyon reach.
- Recent declines in sturgeon recruitment correlate with changes in the biota within the Lower Granite pool.
- We believe trap and transport (upstream and downstream) should benefit this population.
- Flows that now result in "good" recruitment have occurred twice in the last 25 years. If this
 continues, the only way to rebuild and maintain this population may be through a hatchery
 program.






White Sturgeon Population Estimate in the Hells Canyon Reach




What portion of 2017 Year Class entrained?


- 293 fish PIT tagged in 2017-2018 (96% were tagged in 2018)
- 14 were detected being entrained downstream
- 20% of the Acoustic tagged fish that we know entrained also were detected by PIT arrays at the Dam
- 14/20% = 70
- 70/293 = 24% (after two runoff periods)

White Sturgeon Population Estimates




White Sturgeon Survival in Lower Granite Pool







Flows Out of Lower Granite Dam



Flows Out of Lower Granite Dam



Sturgeon Pit Tag Detection and Lower Granite and Little Goose Dams



Flows Out of Lower Granite Dam

